

The APM cluster-galaxy cross-correlation function : Constraints on Ω and galaxy bias.

Rupert A. C. Croft¹, Gavin B. Dalton² & George Efstathiou³

¹*Department of Astronomy, The Ohio State University, Columbus, Ohio 43210, USA.*

²*Department of Physics, University of Oxford, Keble Road, Oxford, OX1 3RH, UK.*

³*Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA, UK.*

1 February 2008

ABSTRACT

We estimate the cluster-galaxy cross-correlation function (ξ_{cg}), from the APM galaxy and galaxy cluster surveys. We obtain estimates both in real space from the inversion of projected statistics and in redshift space using the galaxy and cluster redshift samples. The amplitude of ξ_{cg} is found to be almost independent of cluster richness. At large separations, $r \gtrsim 5 h^{-1}\text{Mpc}$ ($h = H_0/100\text{km s}^{-1}$, where H_0 is the Hubble constant), ξ_{cg} has a similar shape to the galaxy-galaxy and cluster-cluster autocorrelation functions. ξ_{cg} in redshift space can be related to the real space ξ_{cg} by convolution with an appropriate velocity field model. Here we apply a spherical collapse model, which we have tested against N-body simulations, finding that it provides a surprisingly accurate description of the averaged infall velocity of matter into galaxy clusters. We use this model to estimate β ($\beta = \Omega^{0.6}/b$ where b is the linear bias parameter) and find that it tends to overestimate the true result in simulations by only $\sim 10 - 30\%$. Application to the APM results yields $\beta = 0.43$ with $\beta < 0.87$ at 95% confidence. This measure is complementary to the estimates made of the density parameter from larger scale bulk flows and from the virialised regions of clusters on smaller scales. We also compare the APM ξ_{cg} and galaxy autocorrelations directly to the mass correlation and cluster-mass correlations in COBE normalised simulations of popular cosmological models and derive two independent estimates of the galaxy biasing expected as a function of scale. This reveals that both low density and critical density cold dark matter (CDM) models require anti-biasing by a factor ~ 2 on scales $r \leq 2 h^{-1}\text{Mpc}$ and that the Mixed Dark Matter (MDM) model is consistent with a constant biasing factor on all scales. The critical density CDM model also suffers from the usual deficit of power on large scales ($r \gtrsim 20 h^{-1}\text{Mpc}$). We use the velocity fields predicted from the different models to distort the APM real space cross-correlation function. Comparison with the APM redshift space ξ_{cg} yields an estimate of the value of $\Omega^{0.6}$ needed in each model. We find that only the low Ω model is fully consistent with observations, with MDM marginally excluded at the $\sim 2\sigma$ level.

Key words: Galaxies : Clustering ; Large-scale structure of the Universe ; Clusters of galaxies; Cosmology.

1 INTRODUCTION

The spatial cluster-galaxy cross-correlation function $\xi_{cg}(r)$ is defined so that the probability dP of finding a galaxy in the volume element dV at a distance r from a cluster is

$$dP = \bar{n}[1 + \xi_{cg}(r)]dV, \quad (1)$$

where $\bar{n}$ is the mean space density of galaxies. $\xi_{cg}(r)$ is therefore equivalent to the radially averaged overdensity profile of galaxies centred on a typical cluster of galaxies.

The first measurements of galaxy-cluster correlations were made by Seldner & Peebles (1977) who measured the angular cross-correlations of Lick galaxies around Abell clusters. Their results suggested that $\xi_{cg}(r)$ was positive and significantly different from zero out to large spatial separations of $r \sim 100 h^{-1}\text{Mpc}$. A more recent analysis by Lilje & Efstathiou (1988) used cluster redshifts to determine the cross-correlation between Abell clusters and Lick galaxies as a function of metric separation ($w(\sigma)$, see Section 2.2). They found no convincing evidence for clustering on scales

$r \gtrsim 20 h^{-1}\text{Mpc}$ and concluded that some of the signal seen by Seldner and Peebles was due to artificial gradients in the Lick catalogue. However, Lilje and Efstathiou did find some evidence for more large-scale power than predicted by the ‘standard’ (*i.e.* $\Omega = 1$, $h = 0.5$, scale-invariant) CDM model. On scales smaller than $r \lesssim 10 h^{-1}\text{Mpc}$, the real-space $\xi_{cg}(r)$ recovered by inverting the observed form of $w(\sigma)$ is well fit by a power law:

$$\xi_{cg}(r) = \left(\frac{r_0}{r}\right)^\gamma, \quad (2)$$

with $\gamma = 2.2$ and $r_0 = 8.8 h^{-1}\text{Mpc}$.

Direct measurements of the spatial cross-correlation function from redshift surveys of galaxies and clusters (Dalton, 1992, Efstathiou 1993, Mo, Peacock and Xia, 1993, Moore et al. 1994) have confirmed that ξ_{cg} has a similar shape to the galaxy-galaxy and cluster-cluster correlation functions, but with an amplitude roughly equal to their geometric mean. The analyses of Dalton (1992) and Efstathiou (1993) were performed using the original APM cluster sample of Dalton et al. (1992) and the Stromlo-APM galaxies of Loveday et al. (1992).

In this paper we use compute ξ_{cg} using the Stromlo-APM galaxy redshift survey and the sample of 364 clusters of Dalton et al. (1994b). We investigate its behaviour as a function of cluster richness and compare it with the predictions of popular cosmological models. We also calculate $\xi_{cg}(\sigma, \pi)$, the cross-correlation as a function of separation along and perpendicular to the line of sight. The peculiar velocity field around clusters influences the shape of $\xi_{cg}(\sigma, \pi)$ and is expected to depend on Ω , and the mass to light ratio around clusters, so that we can extract some information about the density parameter and galaxy biasing from our measurements. We also carry out some specific comparisons with N-body simulations (so including non-linear effects) to investigate how the biasing of galaxies is expected to vary as a function of scale.

2 CLUSTER-GALAXY CORRELATIONS IN THE APM SURVEY

2.1 The APM data samples

In this paper we will use three different data samples, the APM angular galaxy catalogue, the APM-Stromlo galaxy redshift survey, and the APM cluster redshift survey. The APM galaxy survey (Maddox et al. 1990a, Maddox et al. 1990b, Maddox, Efstathiou and Sutherland 1996) consists of angular positions and other information, but not redshifts, for over 2 million galaxies with a b_J magnitude limit of 20.5. The Stromlo-APM redshift survey (Loveday 1990, Loveday et al. 1992a, Loveday et al. 1992b) is a survey of 1787 galaxies randomly sampled at a rate of 1 in 20 from all APM galaxies with b_J brighter than 17.15. To construct the APM cluster sample, an automated procedure was used to select clusters from the angular APM survey (Dalton 1992, Dalton et al. 1997). Cluster redshifts were then measured, and used to construct an original redshift catalogue of 190 clusters (Dalton et al. 1992) and an extension to 364 clusters (Dalton et al. 1994). It is the latter cluster catalogue which we will use in this paper. In our analyses we will use angular clustering and its inversion to obtain real space 3

dimensional information, using the parent APM galaxy survey and the cluster redshift survey (Section 2.2). We will also measure the 3 dimensional clustering directly from the APM-Stromlo redshift survey and the cluster redshift survey (Section 2.3).

2.2 The projected cross-correlation function

The simplest statistic which can be used to constrain $\xi_{cg}(r)$ is the angular cross-correlation function, $w_{cg}(\theta)$. However, the inversion of this quantity to find $\xi_{cg}(r)$ tends to be rather unstable. Since we are using the APM cluster redshift survey as our cluster sample we can make use of the cluster redshift information to determine the projected cross-correlation function, $w_{cg}(\sigma)$, as defined by Lilje & Efstathiou (1988), where $\sigma = cz\theta/H_0$ is the metric separation of a cluster with redshift z and a galaxy at angular distance θ from the cluster centre. As our estimator for $w_{cg}(\sigma)$ we use the standard estimator,

$$w_{cg}(\sigma) = \frac{N_{ran}}{N_{gal}} \frac{N_{CG}(\sigma)}{N_{CR}(\sigma)} - 1, \quad (3)$$

where CG and CR denote cluster-galaxy and cluster-random pairs, respectively. We account for the window function of the APM Galaxy Survey on a plate-by-plate basis by generating a random catalogue for a single plate with 100000 random points and excising the regions masked from the survey as the catalogue is used for each plate in turn. With this method, we can use individual galaxy positions from the survey data rather than binned cell counts and so we can measure $w_{cg}(\sigma)$ accurately at small scales. On larger scales we have checked that this method does not introduce large-scale power into our determination of $w_{cg}(\sigma)$ by comparing with the results obtained using cell counts for galaxies and a single random catalogue for the whole survey (Dalton 1992). A similar method has been used more recently in the analysis of the Durham-UKST galaxy redshift survey (Ratcliffe et al., 1997).

The data for $w_{cg}(\sigma)$ are shown in Figure 1 for the cluster sample cross-correlated with all galaxies to three different magnitude limits. The error bars shown are obtained by dividing the APM survey region into four quadrants and determining the error on the mean from the scatter between the four zones. Given the estimate of the depth of the Lick catalogue obtained by Maddox et al. (1990a), we would expect the points for $m_{lim} = 18.5$ to correspond to the results for $w_{cg}(\sigma)$ obtained by Lilje & Efstathiou (1988). A comparison of Figure 1 to Figure 9c of that paper reveals good agreement over the range of σ for which w_{cg} can be measured reliably.

The inversion of projected clustering information in the form of $w_{cg}(\sigma)$ to the three-dimensional statistic $\xi_{cg}(r)$ involves a weighted summation of the $w_{cg}(\sigma)$ points (Saunders, Rowan-Robinson & Lawrence, 1992):

$$\xi_{cg}(r) = \frac{-1}{\pi B} \sum_{j \geq i} \frac{w(\sigma_{j+1}) - w(\sigma_j)}{\sigma_{j+1} - \sigma_j} \ln \left(\frac{\sigma_{j+1} + \sqrt{\sigma_{j+1}^2 - \sigma_i^2}}{\sigma_j + \sqrt{\sigma_j^2 - \sigma_i^2}} \right) \quad (4)$$

where $r = \sigma_i$. The factor B in Equation 4 accounts for the difference in the selection functions of the clusters and galaxies and is defined as follows (Lilje & Efstathiou 1988):




Figure 1. The projected cluster-galaxy cross-correlation function for APM galaxies and APM clusters. Results are shown using APM galaxies with different magnitude limits as indicated in the figure. The error bars are determined from the scatter in the results derived from four nearly equal area zones of the APM survey.




Figure 2. The real space cluster-galaxy cross-correlation function for APM galaxies and APM clusters. Results are shown for different magnitude limits, and the error bars have been calculated as described in the caption for Figure 1. For clarity, the $m_{lim} = 18.5$ and $m_{lim} = 20.5$ points have been slightly offset in the r direction. We also show a fit to the data with form given by Equation 6 and the power law (Equation 2) which Lilje & Efstathiou find is a good fit to the cross-correlation of Abell clusters and Lick counts.

$$B = \frac{\sum_i \psi(y_i)}{\sum_i (1/y_i) \int_0^\infty \psi(x) x^2 dx}. \quad (5)$$

Here ψ is the selection function of the galaxy survey and y_i is the redshift of cluster i . The selection function ψ was evaluated using the luminosity function parameters obtained from the Stromlo-APM survey by Loveday et al. (1992). Again we show error bars obtained by inverting the $w_{cg}(\sigma)$ estimates from four quadrants of the survey for each magnitude range. The data show excellent agreement between the three different magnitude limited galaxy samples used, but are not well represented by a single power law.

For the velocity field analysis in Section 4, we will use a fit to the real space APM cross-correlation function. We choose to fit an arbitrary function which is the sum of an exponentially truncated power law and the linear theory correlation function shape of a scale-invariant CDM model with $\Gamma = \Omega h = 0.2$ (denoted as $\xi_{\Gamma=0.2}(r)$ below) normalised so that $\sigma_8 = 1^*$:

$$\xi_{cg}(r) = \left(\frac{r_c}{r}\right)^\gamma e^{\beta r} e^{\eta/r} + \alpha \xi_{\Gamma=0.2}(r). \quad (6)$$

The parameter combination $r_c = 11.7$, $\gamma = 2.6$, $\eta = 0.6$, $\beta = 1.7$ and $\alpha = 1.8$ gives a reasonable fit to the data for all magnitude bins and is plotted as a solid line on Figure 2. A fit is necessary because the noise level in the real space cross-correlation function becomes rather large for $r \gtrsim 10 h^{-1} \text{Mpc}$. The shape of the fit on these large scales is motivated by the shape of the redshift space ξ_{cg} (Section 2.2). We also plot on the same figure the power-law fit (given by Equation 2) which Lilje and Efstathiou (1988) find is a good approximation to the real space cross-correlation function derived from Abell clusters and Lick counts. We can see that on large scales, there is no evidence for any significant excess of power over this fit. The APM results therefore support the conclusions of Lilje and Efstathiou (1988) summarized in Section 1. Our APM results are in agreement with the cross-correlation of APM clusters and Edinburgh-Durham Sky Survey galaxies carried out by Merchán et al. (1997), although their estimated errors are large. It is useful to note that on small scales, $r \lesssim 2 h^{-1} \text{Mpc}$, our error bars on $\xi_{cg}(r)$ are very small, so that we will be able to draw some interesting conclusions about galaxy biasing on small scales from a comparison with theoretical models (Section 5).

2.3 The redshift space cross-correlation function

We estimate $\xi_{cg}(s)$, where s represents the separation of cluster-galaxy pairs in redshift space, using the APM sample of 364 clusters of Dalton et al. (1994b) and the ~ 2000 galaxies with redshifts from the APM-Stromlo bright galaxy redshift survey (Loveday et al. 1992a). The calculation of $\xi_{cg}(s)$ differs from the evaluation of the cluster-cluster correlation function (Dalton et al. 1994b) as the galaxy selection function falls very steeply with increasing distance. Weights must be therefore be applied to the galaxy-cluster pairs to

* Where σ_8 denotes the *rms* amplitude of the mass fluctuations in spheres of radius $8 h^{-1} \text{Mpc}$.

recover the minimum variance estimate of $\xi_{cg}(s)$. This optimal weighting (at least on scales for which $\xi_{cg}(s) \leq 1$) can be shown to be (see Efstathiou 1988, Loveday 1990)

$$w_{ij} \simeq 1/[1 + 4\pi n(r_i)J_3^{cg}(s_{ij})], \quad (7)$$

where

$$J_3^{cg}(s_{ij}) = \int_0^{s_{ij}} \xi_{cg}(x)x^2 dx, \quad (8)$$

r_i is the distance from the observer to galaxy i and s_{ij} the separation of galaxy i and cluster j . To use the formula we must predict roughly what $J_3^{cg}(s)$ will be - here we use the weighting function resulting from a linear theory CDM power spectrum with $\Gamma = 0.2$ and an amplitude twice that of $\xi_{gg}(s)$ measured for APM-Stromlo galaxies (Loveday et al. 1992). After cross-correlating the two catalogues to find all galaxy-cluster pairs, we cross-correlate the clusters with a catalogue of 100000 random points. This random catalogue has the same boundaries and selection function as the galaxy sample. We then use the standard estimator to find $\xi_{cg}(s)$:

$$\xi_{cg}(s) = \frac{N_{ran}}{N_{gal}} \frac{N_{CG}(s)}{N_{CR}(s)} - 1, \quad (9)$$

where $N_{CG}(s)$ and $N_{CR}(s)$ are the galaxy-cluster and cluster-random pairs in the bin interval centred on s and each are weighted using w_{ij} from Equation 7.

Results for the 364 $\mathcal{R} = 50$ clusters are shown as triangles in Fig. 3. It can be seen that there is some curvature in the plot, with a definite break at $s \sim 30 - 50 h^{-1}\text{Mpc}$. The curve crosses $\xi_{cg}(s) = 1$ at roughly $9 h^{-1}\text{Mpc}$, which is intermediate between the behaviour of the galaxy auto-correlation function, $\xi_{gg}(s)$ ($s_0 \simeq 5 h^{-1}\text{Mpc}$, in the notation of Equation 2, Loveday et al. 1992) and the cluster auto-correlation function $\xi_{cc}(s)$ ($s_0 \simeq 14 h^{-1}\text{Mpc}$, Dalton et al. 1994). The error bars, calculated from Poisson statistics and the number of cluster-galaxy pairs in each bin, are relatively small, indicating that $\xi_{cg}(s)$ will be an interesting statistic to compare with theoretical models. We can compare the results for $\mathcal{R} = 50$ clusters with the solid line in this figure, which is the fit to the real space cross-correlation function (Equation 6). For the moment, we will note that $\xi_{cg}(s)$ is marginally higher than $\xi_{cg}(r)$ on large scales, boosted by streaming motions (Kaiser 1987) and smaller for separations less than $\sim 4 h^{-1}\text{Mpc}$ due to the effect of the cluster velocity dispersion. Uncertainties in the cluster redshifts probably play a part in depressing the amplitude of $\xi_{cg}(s)$ on small scales, as for many clusters we have redshifts for only 2 or 3 galaxies (each with their own measurement errors). The error in the cluster centre of mass velocity could, therefore, be as much as a few hundred km s^{-1} . Dalton et al. (1994a) have compared results for APM clusters with many (> 10) measured galaxy redshifts to the redshifts of the brightest galaxy in each cluster. The rms scatter between the two values is 512 km s^{-1} , which should be higher than the error on our cluster redshifts as we use ≥ 2 galaxy redshifts per cluster.

Moore et al. (1994) have found that ξ_{cg} for IRAS galaxies and Abell clusters is insensitive to cluster richness. We have repeated this type of analysis using our APM samples and the results are plotted in Figure 3. We have increased the lower richness cutoff from $\mathcal{R} = 50$ (the full sample) up to $\mathcal{R} = 80$, but as the results show there is no detectable




Figure 3. The spatial cluster-galaxy cross-correlation function for APM-Stromlo galaxies and samples of APM clusters with different lower richness limits. All cluster subsamples are drawn from the sample of 364 clusters (APM $\mathcal{R} \geq 50$), so that 243, 113 and 60 clusters are used in the calculation of the curves for $\mathcal{R} \geq 60$, $\mathcal{R} \geq 70$ and $\mathcal{R} \geq 80$ clusters respectively. The subsamples have mean intercluster separations of $36 h^{-1}\text{Mpc}$ ($\mathcal{R} \geq 60$), $48 h^{-1}\text{Mpc}$ ($\mathcal{R} \geq 70$) and $59 h^{-1}\text{Mpc}$ ($\mathcal{R} \geq 80$). The solid line is a fit (with form given by Equation 6) to the real space ξ_{cg} results for the APM $\mathcal{R} \geq 50$ clusters.

change in the shape or amplitude of $\xi_{cg}(s)$, given the errors. Our different samples have space densities $\bar{n}_c = 3.5 \times 10^{-5} h^3\text{Mpc}^{-3}$ ($\mathcal{R} = 50$), $2.2 \times 10^{-5} h^3\text{Mpc}^{-3}$ ($\mathcal{R} = 60$), $9.3 \times 10^{-6} h^3\text{Mpc}^{-3}$ ($\mathcal{R} = 70$) and $4.7 \times 10^{-6} h^3\text{Mpc}^{-3}$ ($\mathcal{R} = 80$). We might expect the cross-correlation function of richer clusters to have a higher amplitude, at least on small scales, as cluster richness should be related to ξ_{cg} within the cluster selection radius. This was discussed by Seldner and Peebles (1977), who found that the effect was smaller than expected for Abell clusters. This is probably because the distance indicator used by Abell depends on cluster richness; for clusters at a given apparent distance, the richer objects would actually be further away, thus depressing the amplitude of the angular cross-correlation function. These problems should not affect the APM cluster sample as it has been designed so that cluster richness does not affect apparent distance (Dalton et al. 1997). However, the situation here is complicated by the fact that we are measuring redshift space clustering. Rich clusters will have their clustering signal smeared out due to their high velocity dispersions, so that the amplitude of clustering will be more heavily depressed on small scales than for poorer clusters. The underlying situation on small scales is therefore not entirely obvious. On larger scales ($r \gtrsim 1 h^{-1}\text{Mpc}$) though, our results show that the amplitude of $\xi_{cg}(s)$ really does depend only very weakly on cluster richness. We will show in the next section that this is consistent with model predictions.

We have calculated ξ_{cg} for the APM sample in redshift space as a function of pair separation along the line of sight (π) and perpendicular to the line of sight (σ). The effects of peculiar velocities, which distort the pair separations in the π direction are evident in the results for our sample plotted in Figure 4. We can see elongation present on small scales




Figure 4. The APM cluster-galaxy cross-correlation function $\xi_{cg}(\sigma, \pi)$ (calculated using the full sample of 364 clusters) shown as a function of pair separations perpendicular to the line of sight (σ) and along the line of sight (π). Contour levels are at $\xi_{cg} = 4, 3, 2, 1, 0.8, 0.6, 0.4, 0.2, 0.1, 0., -0.05$. The contour level at $\xi_{cg} = 1$ is shown by the heavy line; negative contours are plotted as dashed lines. $\xi_{cg}(\sigma, \pi)$ was calculated using 16 bins in σ and π in the interval $0 - 40 h^{-1}\text{Mpc}$. For clarity, before plotting the results in this figure they were smoothed using a moving window average (3×3 bins).

which is caused by random galaxy velocities and redshift measurement errors. On larger scales, we can see a break in the contours around $\sigma \simeq 6 h^{-1}\text{Mpc}$, $\pi \simeq 10 h^{-1}\text{Mpc}$ which could be caused by the infall region around the cluster. This coherent infall should cause compression of the contours along the σ axis on larger scales (Kaiser 1987, Lilje & Efstathiou 1989), but there does not seem to be obvious evidence of this in Figure 4.

The velocity field around clusters should be dependent on the mass distribution, and therefore on the value of Ω . If we have a simple model for how the two main effects present in velocity field arise (small scale dispersion and large scale infall) we should be able to use distortions in $\xi_{cg}(\sigma, \pi)$ to derive information on Ω and the amplitude of mass fluctuations. To do this, we need to know ξ_{cg} in real space and to have a model which describes the behaviour of galaxy velocities around clusters. We apply both of these to our $\xi_{cg}(\sigma, \pi)$ data from the APM survey in Section 4.3 below. We will first examine the predictions of theoretical models using N-body simulations and use them to develop a model of the velocity field around clusters.

3 ξ_{CG} FROM SIMULATIONS OF COSMOLOGICAL MODELS

3.1 Clusters and galaxies in simulations

Our predictions of the form of the cluster-galaxy correlation function in cosmological models come from studying N-body simulations. We use catalogues of clusters constructed from simulations which have been used in previ-

ous papers to study the cluster-cluster correlation function (Croft & Efstathiou 1994, Dalton et al. 1994b). The simulations are of three different spatially flat universes, two CDM models and one Mixed Dark Matter (MDM) model. One set of simulations is of “standard” CDM (SCDM), with $\Omega_0 = 1, h = 0.5$ and the other is of low density CDM (LCDM) with $\Omega_0 = 0.2, h = 1$ and a cosmological constant Λ , where $\Lambda = 0.8 \times 3H_0^2$. The power spectra for the SCDM and LCDM models are as given Efstathiou, Bond & White (1992). For the MDM model, we used the form given in Klypin et al. (1993) with $\Omega_0 = 1, h = 0.5$, and a massive neutrino component contributing $\Omega_\nu = 0.3$. We assume scale invariant primordial fluctuations for all models.

Each simulation contains 10^6 particles in a box of co-moving side-length 30000km s^{-1} and was run using a P³M N-body code (Efstathiou et al. 1985). We use 5 realisations of each model with different random phases.

In this Section we use simulated cluster catalogues constructed from the N-body simulations to have the same mean separation as the APM sample ($d_c = 30 h^{-1}\text{Mpc}$). The clusters are identified from simulations using a friends-of-friends algorithm to select candidate centres. The mass enclosed within a fixed radius (in this case $0.5 h^{-1}\text{Mpc}$) of the centre of mass is computed and the clusters are ordered by mass. Finally, a mass limit is applied to generate a cluster catalogue of a specified mean space density. The procedure is described in more detail in Croft & Efstathiou (1994). We note here that as long as the clusters are defined to be collapsed objects, the set of objects identified in the simulations is insensitive to the selection criterion. For example, Gaztañaga, Croft & Dalton (1995) identify clusters as high peaks in the density field smoothed with small filters and recover essentially the same catalogue of clusters (for a variety of filter sizes) as our percolation algorithm. We can also reasonably expect the positions and mass rankings of galaxy clusters in our simulations to be insensitive to the details of the galaxy formation process, which in the real Universe would turn a large agglomeration of dark matter into a galaxy cluster. These properties of galaxy clusters make them especially useful for elucidating the nature of galaxy biasing and galaxy peculiar velocities. In essence they consist of a set of fixed reference points around which the galaxy overdensity profiles and galaxy velocity profiles can be compared directly with observations.

We do not attempt to carry out any sort of direct identification of galaxies in the simulations. Instead, we calculate the cluster-particle correlation function, ξ_{cp} . The APM ξ_{cg} observations can be compared to ξ_{cp} to determine what sort of galaxy biasing is needed in each model. According to linear perturbation theory, the amplitude of σ_8 grows in proportion to the growth rate $D(t)$ of linear density perturbations (see Peebles 1980, Section 10). The amplitude of the two-point correlation function of the mass fluctuations thus grows as $\sigma_8^2 \propto D^2(t)$ on scales on which linear theory is applicable. On the other hand, the rich cluster two-point correlation function, is almost independent of time (see Croft & Efstathiou 1994) because clusters are rare objects and are strongly biased compared to the mass distribution. We therefore expect the linear theory growth rate of ξ_{cp} to lie between these two cases, and to be proportional to σ_8 , as in the high-peak model of Bardeen et al. 1986. This means that the choice of output time (and hence normal-




Figure 5. The cluster-mass cross-correlation function for simulations of (a) Standard CDM, (b) Low Density CDM and (c) Mixed Dark Matter (see text for the parameters of these models). Results are plotted in real space (solid lines) and in redshift space (dashed lines). All models are normalised to be consistent with the amplitude of fluctuations measured by COBE (ignoring tensor modes), so that $\sigma_8 = 1.0$ for panels (a) and (b) and $\sigma_8 = 0.67$ in panel (c). The model curves in panel (c) have been scaled upwards by a biasing factor of 1.5 in order to make the model consistent with the amplitude of galaxy fluctuations. Also plotted in each panel (as triangles) is the cross-correlation function of APM Stromlo galaxies and the APM cluster sample described in Section 2.3. The error bars on these points have been calculated using Poisson statistics. The dotted lines show a fit (Equation 6) to the real space cluster-galaxy correlation function, consistent with measurements from the APM survey and also with the cross-correlation of Abell clusters and Lick counts.

isation of the model) will make some difference to our results. The normalisation we have used is consistent with the amplitude of fluctuations inferred from COBE microwave background temperature anisotropies (Wright et al. 1994) so that $\sigma_8 = 1.0$ for SCDM and LCDM and $\sigma_8 = 0.67$ for MDM. If we assume that σ_8 for APM galaxies is close to 1.0 (Gaztañaga 1995) and density evolution on this scale has been linear, then ξ_{cg} for SCDM and LCDM should be very nearly equal to ξ_{cg} . In the case of MDM, when plotting our results we merely scale the curve upwards by a constant linear biasing factor, $1.0/\sigma_8$. A discussion of more complicated biasing, including variations with spatial scale which might be required in some models, is deferred to Section 5.

3.2 Results

Our results for these three models are shown in Figure 5, both in real space (solid lines) and redshift space (dashed lines). Concentrating on the real space results first, it can be seen that $\xi_{cg}(r)$ exhibits a sudden change of slope on small scales in all cases. The SCDM and LCDM models have the steepest slopes for this part of the curve, which we might expect, given that they have the most small scale power. The redshift space results show the usual depression on small scales and amplification on large scales, both effects being largest in the case of the two $\Omega = 1$ models, which have the largest particle velocities. As far as a comparison with the APM $\xi_{cg}(r)$ is concerned, we can see that the usual discrepancy on large scales with SCDM is evident, but that the shape on these scales is consistent with LCDM. Indeed for $r > 3 h^{-1}\text{Mpc}$ LCDM appears to give a good fit in both

real and redshift space. On smaller scales, MDM appears to give a reasonable fit to the shape of both the real space results (Equation 6) and those in redshift space. We should be cautious about drawing firm conclusions from this, as the MDM model has been simulated without including the thermal velocities of massive neutrinos and this may affect the density profiles and internal structure of the clusters. The linearly biased MDM model does also seem to have a rather high amplitude on intermediate scales, particularly in redshift space (a point that we will return to in Sections 4 and 5). In any case, it is possible that the efficiency of galaxy formation is different near clusters leading to scale dependent biasing. We will investigate this possibility in Section 5.

The richness dependence of ξ_{cg} in the models is shown in Figure 6, where we plot $\xi_{cg}(s)$ (in redshift space) for simulated clusters with a similar range of space densities to the 4 different APM samples shown in Figure 3. There is a very weak richness dependence in the amplitude of ξ_{cg} , which becomes even weaker on large scales ($s \geq 25 h^{-1}\text{Mpc}$). This weak dependence of ξ_{cg} with cluster richness is compatible with the observational results for the APM samples presented in Figure 3, which show no significant dependence of ξ_{cg} with cluster richness.

4 REDSHIFT SPACE DISTORTIONS AND $\xi_{CG}(\sigma, \pi)$




Figure 6. The cluster-mass cross-correlation function in redshift space for clusters with different lower mass limits in simulations of (a) Standard CDM, (b) Low density CDM and (c) Mixed Dark Matter. The curves are labelled with the mean intercluster separation of each sample.

4.1 The spherical infall model

There exists a regime between streaming of galaxies on large-scales and the virialised region of clusters which is important in modelling observations of $\xi_{cg}(\sigma, \pi)$. As the density enhancement within a few $h^{-1}\text{Mpc}$ of clusters is greater than unity, linear theory is not expected to describe the velocity field accurately. However, if we assume that clusters are spherically symmetric, a solution for the non-linear collapse of the system can be found which is exact before orbit crossing takes place. The solution is obtained by treating a proto-cluster with a top-hat profile as if it were an isolated Friedmann universe with its own value of Ω_0 (see e.g. Regös & Geller 1989 for details). Here, we use a good approximation to the exact solution due to Yahil (1985) (also used by Lilje & Efstathiou 1989), who gives the following expression:

$$v_{infall}^{non-lin}(r) = -\frac{1}{3}\Omega_0^{0.6}H_0r\frac{\delta(r)}{(1+\delta(r))^{0.25}}, \quad (10)$$

where $\delta(r)$ is the overdensity inside radius r ,

$$\delta(r) = \frac{3J_3^{cp}(r)}{r^3}, \quad J_3^{cp}(r) = \int_0^r \xi_{cp}(x)x^2 dx. \quad (11)$$

By way of comparison, the linear theory prediction of the infall velocity is

$$v_{infall}^{lin}(r) = -\frac{1}{3}\Omega_0^{0.6}H_0r\delta(r). \quad (12)$$

Galaxy surveys provide with information on the overdensity of galaxies inside radius r and not of the mass. Some assumptions about the relationship between galaxies and mass are therefore required to infer a value of Ω_0 from a measurement of $v_{infall}(r)$. Here we use the simple linear biasing picture, so that $J_3^{cp}(r) = J_3^{cg}(r)/b$. We therefore parametrise our infall model using $\beta = \Omega_0^{0.6}/b$. As even the non-linear velocity model is not expected to hold in the cores of clusters, near and in the virialised region, we choose to truncate the

expression for $v_{infall}^{non-lin}(r)$ with an exponential $e^{-\delta(r)/\delta_{cut}}$, with $\delta_{cut} = 50$. (See the comparisons with N-body simulations in the next subsection for a justification of this value of δ_{cut} .)

One further ingredient in our velocity model is the random velocity dispersion about the smooth infalling flow. To make things as simple as possible, we assume a velocity dispersion independent of distance from the galaxy to the cluster, and independent of direction (whether transverse to the line between galaxy and cluster, or parallel to it, for example). We also assume that the velocities are drawn from a Gaussian distribution. The one-dimensional velocity dispersion will therefore be parametrised by one number, σ_v .

4.2 Direct tests of spherical infall on the velocity field in simulations.

In our analysis we need to assume that the velocity field predicted from the average cluster density profile is equivalent to the average cluster velocity field. This is a non-trivial assumption and must be tested in some way (in our case we will use simulations) before we can make any claims for the reliability of our results. The spherical infall model has previously been used in many studies of the infall regions around individual rich clusters, particularly Virgo (e.g. Yahil, Sandage & Tammann 1980, Yahil 1985). However, the assumption of spherical symmetry for any individual cluster is difficult to justify empirically. In contrast, the average cluster profile measured by $\xi_{cg}(r)$ possesses spherical symmetry by construction. Here we test the validity of our dynamical approximations by comparing our predicted velocity fields directly with the velocity fields measured around N-body clusters. We present results both for the mean infall velocity as a function of radius and the dispersion about the mean infall.

The average infall velocities of particles, are plotted as




Figure 7. The mean infall velocity of particles around simulated clusters as a function of radius. The filled circles in each panel correspond to the infall velocities found in N-body simulations of (a) SCDM, (b) LCDM and (c) MDM. The error bars show the error on the mean calculated from the scatter in an ensemble of 5 realisations. In each panel we have also plotted the predictions of linear theory (dashed lines) and a non-linear spherical infall model (solid lines), as described in the text. The two pairs of dashed and solid lines show the effect of including and not including an exponential truncation at high overdensities as described in Section 4.1.




Figure 8. The one-dimensional velocity dispersion of particles around simulated clusters as a function of radius. The filled circles in each panel correspond to the dispersion in the component of relative velocity along the line from particle to cluster (ie. dispersion about the mean streaming motion plotted in Figure 7). The open circles show the one-dimensional dispersion in the transverse direction. We plot the same models as in previous diagrams: (a) SCDM, (b) LCDM and (c) MDM. The error bars again show the error on the mean calculated from the scatter in an ensemble of 5 realisations.

a function of distance from the cluster centre in Figure 7 (filled circles). We also plot the predictions of spherical infall (solid lines) and linear theory (dashed lines). The rapid decrease of infall velocity at small r in the simulations is due to particles reaching the boundary of the virialised region. We have roughly approximated this effect in our velocity models by the exponential truncation described in Section 4.1. The pairs of dashed and solid curves show the velocity models with and without the exponential truncation. Because of the somewhat arbitrary nature of this truncation, we will restrict our quantitative use of the velocity model to the

region $r > 2.5 h^{-1}\text{Mpc}$. Evidently, the spherical non-linear model provides a much better match to the N-body results than the linear. Over much of the range of density contrasts relevant to our study of the infall region, the non-linear velocity model underpredicts the infall by roughly constant factor, but this is only $\sim 30\%$ for LCDM and $\sim 10 - 20\%$ for SCDM and MDM. At larger radii, $r \geq 20 h^{-1}\text{Mpc}$ we see that both linear and non-linear models begin to match the N-body results even more closely. We therefore expect that when we come to estimate β from distortions in the

cross-correlation function our results will be fairly close to the true value.

The underprediction of the spherical infall model is in agreement with results found by Van Haarlem (1992) and Diaferio and Geller (1996). As the velocity field in the immediate vicinity of galaxy clusters has the potential to be complicated, there have been many mechanisms suggested that might disrupt simple spherical infall. Among these are the presence of shear which should speed up collapse (Hoffman 1986, Lilje & Lahav 1991), which does not appear to be happening here, and formation of substructure and/or ellipticity of the proto-cluster which slows infall (e.g. van Haarlem & van de Weygaert 1993). The study of clusters in the SCDM scenario carried out by Villumsen & Davis (1986) differs from ours in that they analysed the velocity field around individual clusters. Here we are dealing with the averaged velocity around all clusters.

Figure 8 is a plot of the 1-dimensional relative velocity dispersion as a function of distance from particle to cluster in our three cosmological models. The filled symbols show the dispersion about the mean infall motion along the line from particle to cluster and the open circles show the one-dimensional dispersion in the component of relative velocity transverse to this line. It can be seen that our assumption that σ_v is independent of distance and direction turns out to be surprisingly good. The plots also show that the transverse σ_v is slightly higher than the radial value, in the infall region between $r \sim 2 h^{-1}\text{Mpc}$ and $r \sim 10 h^{-1}\text{Mpc}$. This might be the signature of some sort of “previrialisation” occurring (Davis & Peebles 1977, Peebles 1993).

To illustrate the complexity of the velocity field around the simulated clusters, and to see if we can understand why some scenarios fit the spherical infall picture slightly better than others, we have plotted the particle velocities around a sample of individual clusters. In Figures 9 to 11 we show the x and y components of the smoothed peculiar velocity field (a $2 h^{-1}\text{Mpc}$ Gaussian filter was used) around 9 clusters for each model, in the rest frame of the cluster (plotted at the centre of each panel). We have also plotted the density field smoothed with the same filter as a grayscale. We can reach several conclusions from studying these plots:

- The magnitude of the velocity field around MDM clusters (and to a lesser extent SCDM) is anisotropic - there are several clusters with small arrows on one side and large ones on the other.
- The larger coherence in the velocity field of the MDM model compared to the two CDM models is evident. Also obvious is the greater magnitude of the velocities in this model.
- There is substructure in the density field for all models and the cluster we are interested in at the centre of the panel is often part of an elongated system. The MDM model also has a noticeably smoother density field on small scales.
- All models contain some clusters where the main flow pattern is not centered on the cluster but continues past it, and others have more complex and non-radial flows. The pictures do not inspire confidence in the idea of simple spherically symmetric infall around each cluster. It seems that this picture is only likely to apply after some sort of averaging is carried out, at least around each cluster (e.g. as in Villumsen & Davis 1986) or around a statistically defined sample of clusters, as described in this paper. It is also worth consider-

ing that these models do have a relatively high amplitude of mass fluctuations, being normalized to COBE rather than to give the correct abundance of galaxy clusters (White, Efstathiou & Frenk 1993). It is possible that lower amplitude models would conform better to a smooth infall picture.

Obvious signatures of radial infall, such as overdensity “caustics” at the radius of turnaround from the Hubble flow (Regös & Geller 1989) have been difficult to find in observations of real clusters and in N-body simulations (van Haarlem et al. 1993). This does not seem surprising given the complexity evident in these plots. This is consistent with the work of Colberg et al. (1997) who have shown that matter infalling onto individual clusters in simulations does so from specific directions which are correlated with the surrounding large scale structure. It is possible that modelling the velocity and density fields as triaxial ellipsoids (van Haarlem & van de Weygaert 1993) may help in the understanding of the flow. In any case the accuracy of our spherically averaged approximation is good enough to permit a quantitative analysis of the redshift space distortions.

4.3 Estimates of β and σ_v

We are now in the position to make some predictions of how the real space ξ_{cg} will be distorted as a function of β and σ_v . To do this, we shall convolve $\xi_{cg}(r)$ with our velocity model, so that ξ_{cg} in redshift space, $\xi_{cg}(\sigma, \pi)$, is given by (Peebles 1993, Section 20)

$$\xi_{cg}(\sigma, \pi) = (\sqrt{2\pi}\sigma_v)^{-1} \int_{-\infty}^{\infty} [\exp(v - [(v_{infall}^{non-lin}(r)y)/\sigma_v])^2 \times (1 + \xi_{cg}(r)) - \exp(-v^2/\sigma_v^2)] dv, \quad (13)$$

where $r = \sqrt{(\sigma^2 + \pi^2)}$ and $y = \sqrt{(r^2 - \sigma^2)}$. To model the observed estimates of $\xi_{cg}(\sigma, \pi)$, we model $\xi_{cg}(r)$ in Equation 13 with the fitting function of Equation 6 which is consistent with the real space cross-correlation function of the APM sample (Section 2.2). We also test Equation 13 using the N-body simulations and empirical estimates of $\xi_{cg}(r)$ calculated in real space (plotted in Fig. 5).

To place constraints on β and σ_v we calculate the expected $\xi_{cg}(\sigma, \pi)$ for a closely spaced grid of values, over the range $\beta = 0 - 1.5$ and $\sigma_v = 0 - 1500\text{km s}^{-1}$. $\xi_{cg}(\sigma, \pi)$ is calculated at 16×16 points, spanning the range of values for π and σ plotted in Figure 4. This is done by convolving the real space $\xi_{cg}(r)$ with our velocity model using Equation 13. In Figure 12 we show contour plots for a few sample values of β and σ_v . We then find which of our grid of models has the smallest χ^2 by comparing with the observed $\xi_{cg}(\sigma, \pi)$. We estimate confidence bounds on β and σ_v from the distribution of $\Delta\chi^2$ over the $[\beta, \sigma_v]$ plane. As the infall model does not give an accurate prediction for the velocity in the virialised region, we have chosen to exclude from the fit the $\xi_{cg}(\sigma, \pi)$ point with the smallest value of σ and π . Thus we exclude any cluster-galaxy pairs with both σ and $\pi < 2.5 h^{-1}\text{Mpc}$.

The χ^2 contours are shown in Figure 13, where we have tested the SCDM, LCDM and MDM simulations. For the test on the N-body models, we have used all (≈ 1000) clusters in each box (and all 10^6 particles), as well as averaging over 5 realisations. We can see straight away that we are predicting values, for β at least, that are consistent with the results of our direct comparison of infall velocities plotted in Figure 7. For the LCDM model, for which




Figure 9. The distribution of particles (with their projected velocities) surrounding 9 clusters taken from a simulation of a standard CDM universe (with $\sigma_8 = 1.0$). The clusters plotted correspond to those ranked by mass as numbers 2,4,8,16,32,64,128,256 and 512 from a box of size $300 h^{-1}\text{Mpc}$. In each panel, the cluster in question is located in the centre, and the velocity field of particles (in the rest frame of the cluster) around the cluster has been assigned to a grid using a $2 h^{-1}\text{Mpc}$ Gaussian kernel. The X and Y components of this velocity field in a slice through the centre of the cluster are shown as arrows. The particles have also been assigned to a grid using the same kernel and a slice through the resulting density field is also shown on each panel as a grayscale (linear in density).

the true value is $\beta = 0.39$ (Lahav et al. 1991), we find $\beta = 0.52^{+0.06}_{-0.09}$ (95% confidence limits). For SCDM and MDM we measure $\beta = 1.20^{+0.09}_{-0.12}$ and $\beta = 1.14^{+0.12}_{-0.18}$, respectively. The error bars on β given here are computed by marginalisation over all values of σ_v . The confidence limits on the pair of parameters σ_v and β are plotted as contours in Figure 13. This diagram shows that given enough clusters we should be able to distinguish between high and low β models. This constraint on β from the infall region around clusters (most of the signal comes from $r < 10 h^{-1}\text{Mpc}$) is complementary both to measurements of the central mass of clusters from the virial theorem and to larger scale bulk flow observations (e.g. Loveday et al. 1996). As the infall method tends to systematically overestimate β by a small amount, any measurement which gives a particularly low value, for example one inconsistent with high Ω models, will be especially interesting.

The values of σ_v we measure using the maximum likelihood fits are $370^{+60}_{-90}\text{km s}^{-1}$, $690^{+60}_{-60}\text{km s}^{-1}$, and $570^{+30}_{-60}\text{km s}^{-1}$ for LCDM, SCDM and MDM respectively. These are again 95% confidence limits obtained after marginalising over all values of β . Comparison with Fig. 8

reveals that these values are in good agreement with the velocity dispersion of particles in the outer parts of clusters.

We now turn to the determination of β and σ_v from the APM sample. The results of the maximum likelihood fitting are shown in panel (d) of Figure 13. We can see immediately that we have much larger statistical errors than in our simulations, as there are only 364 clusters and ~ 2000 galaxies in the APM sample. The value of β we obtain is 0.43, with $\beta < 0.87$ at 95% confidence. The tail of probabilities towards rather high values of σ_v and the best fit σ_v inferred, 660km s^{-1} ($\sigma_v < 1070\text{km s}^{-1}$ at 95% confidence) are probably a consequence of uncertainties in the cluster redshifts. There is not therefore a good constraint on the velocity dispersion in the outer regions of clusters from the APM sample.

5 MODEL-DEPENDENT COMPARISONS: BIASING AS A FUNCTION OF SCALE.

To calculate how biasing of galaxy fluctuations is expected to change with scale for each cosmological model, we calculate




Figure 10. As for Fig 9 but for an LCDM simulation (with $\sigma_8 = 1.0$).

the particle-particle two point correlation function, $\xi_{\rho\rho}(r)$ as well as the cluster-particle cross-correlation function $\xi_{c\rho}(r)$. Given the observational results for $\xi_{gg}(r)$ and $\xi_{cg}(r)$, we can define two biasing relations for each cosmological model,

$$b_{gg}^2(r) = \xi_{gg}(r)/\xi_{\rho\rho}(r) \quad (14)$$

and

$$b_{cg}(r) = \xi_{cg}(r)/\xi_{c\rho}(r). \quad (15)$$

If the efficiency of galaxy formation is not changed by proximity to clusters, then these quantities, $b_{gg}(r)$ and $b_{cg}(r)$, will be equal. We use the estimates of $\xi_{gg}(s)$ calculated from the APM-Stromlo redshift survey of Loveday et al. (1992) in Equation 14. As the observed estimates of ξ_{gg} and ξ_{cg} are computed in redshift-space, we compare them with redshift-space estimates of $\xi_{\rho\rho}$ and $\xi_{c\rho}$ computed from the N-body simulations. The results for the three different models are shown in Figure 14. In each panel we show both $b_{gg}(s)$ and $b_{cg}(s)$. It is clear that for SCDM, the required bias must increase strongly towards larger scales. For LCDM, the derived b_{gg} is roughly equal to unity and nearly independent of scale. There does appear to be a discrepancy between $b_{cg}(s)$ and $b_{gg}(s)$ on small scales, but it is likely that errors in the cluster redshifts have suppressed the small-scale amplitude of $\xi_{cg}(s)$ leading to an underestimate of b_{cg} . The MDM model requires scale-dependent biasing, at least on scales

$\lesssim 5 h^{-1}\text{Mpc}$. The dashed lines in the figures show the expectations of linear theory and constant biasing for models normalised to the COBE measurements according to the linear theory value of σ_8 . In the case of LCDM, this agrees well with the non-linear simulations, although for MDM we can see that non-linear evolution appears have made the matter fluctuation amplitude higher, and less biasing is necessary than linear theory predicts.

On smaller scales ($s \leq 4 h^{-1}\text{Mpc}$), we have seen that $b_{cg}(s)$ appears to be systematically lower than $b_{gg}(s)$ for LCDM and, to a lesser extent, for MDM. As mentioned above, errors in cluster redshifts could be causing this effect. Furthermore, the relation between $b(s)$ in redshift-space and the underlying $b(r)$ depends on the velocity field in each model and so mixes physical scales in a complex way. We will therefore turn to the real space information that we have available, in the form of $\xi_{cg}(r)$ estimated by inversion of the projected APM cluster-galaxy cross-correlation function (Section 2.2). Recovery of spatial correlation functions from projected and angular statistics can depend sensitively on the luminosity function assumed. In Figure 15 we plot $b_{cg}(r)$ obtained when three different magnitude limits are applied to the angular galaxy catalogue (note the difference in the scale of the abscissa compared to Figure 14). On the small scales we are considering here, there is not much difference between the results and we can see that both the CDM mod-




Figure 11. As for Fig 9 but for a MDM simulation (with $\sigma_8 = 0.67$).

els appear to require some anti-biasing. This is in agreement with the study made by Jenkins et al. (1998) who derive an estimate of bias in models as a function of scale in real space from galaxy-galaxy clustering (using Equation 14). Jenkins et al. use a critical density CDM model which has a lower amplitude, and so although that model requires $b_{gg}(r) > 1$ on all scales, there is a relative antibias on small scales compared to large. Our MDM models are reasonably consistent with a constant biasing factor, consistent with the results shown in Figure 5. Of course as no “hot” particles have been used to simulate the neutrinos directly in these models, there is some uncertainty in the accuracy of the cluster profiles on small scales.

It should be added that the models which require anti-biasing do so on scales which are too small to be probed by our non-linear infall model ($r \lesssim 2 h^{-1} \text{Mpc}$). On larger scales, though, redshift distortions of ξ_{cg} can give us more information about biasing in our models. For example, we would expect the bias factors that result from Equations 14 and 15 to be the same when measured in real and redshift space, at least on large scales where linear theory should hold. On examination of Figures 14 and 15, this does appear to be the case for LCDM, but there is the suggestion that the bias factor for both SCDM and MDM is higher in real space than redshift space. This can also be seen from the plot of ξ_{cg} shown directly in Figure 5. This is presumably

because the coherent peculiar velocities in these two models which boost ξ_{cg} in redshift space (Kaiser 1987) are too high. This is just another way of saying that the amplitudes of our models (which are COBE normalised) are probably wrong. Lower amplitudes for these two models, such as those given by using a normalisation which gives the correct space density of galaxy clusters (White, Efstathiou & Frenk 1993, Eke et al. 1996) would presumably match the data better. Changing the normalization would of course not remedy the fact that SCDM does not match the shape of clustering on large scales.

A potentially more powerful way of comparing the models and observations involves using the full $\xi_{cg}(\sigma, \pi)$ information. To do this, we first make the assumption that some form of scale-dependent biasing holds so that $\xi_{cg}(r)$ in the model being tested and the observations are related by Equation 15. We then assume that the velocity field around clusters can be predicted from $\xi_{cp}(r)$ in the model, and that the observed form of $\xi_{cg}(\sigma, \pi)$ has been distorted by this velocity field. We use Equation 13 to model the distortion, with the non-linear infall prediction being computed from $\xi_{cp}(r)$, and used to distort $\xi_{cg}(r)$. At this point, we are guaranteed by construction that the amplitude and shape of the scale-dependent biasing is correct, but using different values of Ω and σ_v in Equation 13 will result in differing degrees of agreement with the observed $\xi_{cg}(\sigma, \pi)$. We carry out χ^2




Figure 12. A smooth fit (Equation 6) to the observed cluster-galaxy cross-correlation function $\xi_{cg}(r)$ convolved with a velocity model (spherical infall and constant velocity dispersion- see text) to produce a predicted $\xi_{cg}(\sigma, \pi)$ plot. We show results for 3 different values of β and 3 different values of the velocity dispersion, σ_v . Contour levels are the same as in Figure 4

fitting of $\xi_{cg}(\sigma, \pi)$ exactly as in Section 4.3, except that we are constraining $\Omega^{0.6}$ required in each specific model, rather than β for a general scenario with scale-independent biasing.

The results of this procedure are shown in Figure 16, where we can see that all models prefer a relatively low value of $\Omega^{0.6}$, which of course is natural for LCDM but not for the others. After marginalising over all values of σ_v , we find the following values for $\Omega^{0.6}$ (with 95% upper limits), 0.49 (1.28), 0.38 (0.88) and 0.32 (1.06), for SCDM, LCDM and MDM respectively. Given the tendency of the spherical infall model to slightly overestimate $\Omega^{0.6}$, these results are reasonably significant. The σ_v results for all models are $\sim 700 \text{ km s}^{-1}$. As before, we can attribute this high value to the large errors in the redshifts of individual APM clusters which should bias the σ_v results upwards. It is interesting that the SCDM model and to a lesser extent the MDM model are expected to give σ_v almost this high without accounting for the uncertainties in the velocities. As we know that these errors are present (Dalton et al. 1994a), we can regard this as additional evidence against these two models.

In this analysis, we have two parameters, σ_v and $\Omega^{0.6}$, each of which probes a different part of the velocity field around clusters. If the model is correct then the best fit values from the observations should match the values of both parameters predicted from the model. We have seen that this fitting procedure favours the LCDM model over the other two we consider here. However, we have been restricted to a narrow set of model parameters and it is possible that by varying these it might be possible to improve the agreement with observations. For example, a lower amplitude SCDM model would probably require a higher value of Ω to fit the data and so be more consistent.

This sort of analysis could also be applied to the galaxy-galaxy correlation function, although the results would probably not be as reliable. The analysis of Section 4 shows that non-linear dynamical model of cluster-galaxy peculiar velocities works accurately even on small scales. However, it has proved difficult to develop a model of galaxy-galaxy velocities of comparable accuracy (see e.g. Hatton and Cole 1997). Throughout our analysis we have neglected the possibility of velocity biasing. Velocity biasing could, if significant, introduce systematic errors in analyses of redshift-space distortions. Unfortunately, theoretical predictions for velocity bias are so uncertain (e.g. Summers, Davis and Evrard 1995) that it is not yet possible to make reliable models of this effect.

6 DISCUSSION AND CONCLUSIONS

The analysis presented in this paper differs from other recent work on the density and velocity fields around clusters in a number of ways. The CNOC cluster redshift survey of Carlberg et al. (1997) specifically targets galaxies within and close to the virialised region of clusters. Carlberg et al. find that a rescaling of cluster profiles allows all clusters to be fit by the universal dimensionless halo profile of Navarro, Frenk and White (1997). We do not carry out such a rescaling here, as we are interested in the profiles of clusters out to a much larger distance (where presumably rescaling would not work, as all profiles must converge to the mean density). Our dynamical measure of the dark matter in each cluster is also based on a different velocity and density regime, the infall region, rather than the virialised region.

The sort of analysis we employ here could be extended




Figure 13. Contours of constant $\Delta\chi^2$ corresponding to 68%, 95% and 99.7% confidence limits on values of σ (the 1 dimensional velocity dispersion) and β ($\simeq \Omega^{0.6}/b$) found by fitting a velocity model to $\xi_{cg}(\sigma, \pi)$. In panels (a) to (c) we show the results found from clusters identified in 5 ensembles each of simulations of the SCDM, LCDM and MDM models respectively. In panel (d) we show results for the APM survey (using the $\xi_{cg}(\sigma, \pi)$ plotted in Figure 4). In each panel the values of σ and β for which the χ^2 was evaluated are shown by dots, and the best fit values are indicated by a cross.




Figure 14. The bias $b(s)$ as a function of scale in redshift space for 3 cosmological models. $b(s)$ is found by dividing the APM $\xi_{cg}(s)$ by $\xi_{cp}(s)$ for each model, the results being represented by open circles in each panel. A separate estimate of $b(s)$ can be found by evaluating $\sqrt{\xi_{gg}(s)/\xi_{pp}(s)}$ and this quantity is plotted as filled circles. The dotted line shows in each case the relation that would be expected if there exists a linear biasing relation between galaxies and mass and σ_8 for galaxies = 1.0.




Figure 15. The bias $b_{cg}(r)$ as a function of scale in real space for 3 cosmological models for the region within $10 h^{-1}\text{Mpc}$ of cluster centres. $b(r)$ is found by dividing the APM $\xi_{cg}(r)$ by $\xi_{cp}(r)$. The real space cross-correlation function is found by inverting the projected cross-correlation function of APM galaxies and clusters (see Section 2.2). Results for three different galaxy magnitude limits are shown, $m_{lim} = 18.5$ (filled circles), $m_{lim} = 19.5$ (open circles) and $m_{lim} = 20.5$ (open squares). The amplitude of the models has been adjusted to our usual COBE normalisation so that $\sigma_8 = 1.0$ for panels (a) and (b) and $\sigma_8 = 0.67$ for panel (c). The dotted line shows in each case the result for $b_{cg}(r)$ that would be expected if there exists a linear biasing relation between galaxies and mass and if σ_8 for galaxies = 1.0.




Figure 16. $\Delta\chi^2$ distributions for $\Omega^{0.6}$ and σ_v evaluated in a similar way to those in Figure 13, but this time using the velocity field predicted from the mass distribution in models to distort the real space ξ_{cg} . This will give an estimate of the values of $\Omega^{0.6}$ and σ_v needed in the models to account for the redshift distortions, given that they are automatically correctly biased to give the same real space ξ_{cg} as the APM survey.

to constrain Ω from the virialised region. In the usual analysis of cluster velocity dispersions, there is no way of telling unambiguously whether a given galaxy is a cluster member with a high velocity or a foreground or background galaxy. Non-member galaxies and the various ways employed to prune them can add significant bias to determinations of the velocity dispersions from samples of clusters (see e.g. van Haarlem et al. 1997). Analysis of the “finger of God” effect on the cross-correlation function offers a way of treat-

ing the problem statistically, which automatically accounts for contamination. In this paper, we have not made use of the information on σ_v because we have a relatively small number of galaxies in our sample and because there are uncertainties in the errors of the APM cluster redshifts. Future surveys such as the 2dF and Sloan redshift surveys (Colless 1997, Gunn & Weinberg 1995) and also the presently available Las Campanas Redshift survey (Schectman et al. 1996) could be used effectively for this purpose. These large sur-

veys will also form good datasets to which we can apply the non-linear infall models used in this paper, and should provide accurate estimates of Ω and biasing parameters from the infall region of clusters.

The value of β we obtain in this paper is consistent with that obtained from a consideration of linear redshift distortions of the galaxy-galaxy clustering function on larger scales (e.g. Loveday et al. 1996). This suggests that any significant biasing as a function of scale is probably confined to scales smaller than $\sim 2 h^{-1}\text{Mpc}$.

In summary, we have investigated the shape and amplitude of cluster profiles both in the APM survey and in simulations of cosmological models with Gaussian initial conditions. In all cases we find a distinct two-component structure, with the shape of $\xi_{cg}(r)$ on large scales consistent with the galaxy-galaxy or particle-particle correlation functions. For $r \lesssim 5 h^{-1}\text{Mpc}$ we see a steepening of the profile. This region is also the part most sensitive to differences in the richnesses of clusters used to calculate $\xi_{cg}(r)$, although the effects are too small to be noticeable in the observational sample.

As expected for objects forming in a bottom-up scenario, the velocity and density field around clusters in the simulations is complex, with substructure evident in plots. Nevertheless, when we average the infall velocities around all clusters, the shape and amplitude of the infall curve is surprisingly well described by a spherically symmetric collapse model. It is therefore feasible to use this simple velocity model to describe distortions in $\xi_{cg}(\sigma, \pi)$ and therefore to constrain Ω . When we do this using the APM data, we find that $\Omega^{0.6}/b$ has a best fit value of 0.43 and is constrained to be less than 0.87 at 95% confidence, where b is the linear bias parameter.

We have also compared ξ_{gg} and ξ_{cg} with N-body simulations of the mass distribution to estimate the biasing of galaxy fluctuations as a function of scale. We conclude that anti-biasing on scales $r \lesssim 2 h^{-1}\text{Mpc}$ is required in the SCDM and LCDM scenarios, as their cluster profiles appear to be steeper than in the observations. This sort of biasing might be possible, as galaxies or quasars forming in clusters might suppress galaxy formation nearby (e.g. Babul & White 1991). On the other hand, this might be hard to reconcile with a galaxy density field which is totally unbiased on large scales (as may be required in LCDM). The MDM model does have the correct cluster profiles, without any scale-dependent biasing. However, we find that the redshift distortions in the cross-correlation function predicted by this model are too strong, by about 2σ , so that the LCDM model should still be considered the most successful of those tested.

Application of the techniques presented in this paper to larger redshift surveys should yield robust measures of the cosmic density from the infall region around galaxy clusters. The spherical infall model is a dynamical approximation which is accurate in the highly non-linear regime and should continue being useful for constraining Ω and biasing in a fashion complementary to traditional linear analyses of galaxy-galaxy clustering.

ACKNOWLEDGMENTS

This work was supported by grants from the UK Particle Physics and Astronomy Research Council, and by NASA Astrophysical Theory Grants NAG5-2864 and NAG5-3111. GPE acknowledges the award of a PPARC Senior Fellowship.

REFERENCES

- Babul, A. & White, S. D. M., 1991, *MNRAS*, **253**, 31P.
- Bardeen J.M., Bond J.R., Kaiser N. & Szalay A., 1986, *ApJ*, **304**, 15.
- Carlberg, R. G., Yee, H. K. C., Ellingson, E., Morris, S. L., Abraham, R., Gravel, P., Pritchet, C. J., Smecker-Hane, T., Hartwick, F.D. A., Hesser, J. E., Hutchings, J. B., Oke, J. B., 1997, *ApJ Lett.*, **485**, L13.
- Colberg, J. M., White, S. D. M., Jenkins, A., & Pearce, F. R., 1997, *MNRAS*, *submitted*, astro-ph/9711040.
- Colless, M., 1997, in *Wide Field Spectroscopy*, eds Kontizas M., Kontizas E., Kluwer.
- Croft, R. A. C. & Efstathiou, G., 1994, *MNRAS*, **267**, 390.
- Dalton, G. B., 1992, *D.Phil thesis*, Oxford University.
- Dalton, G. B., Efstathiou, G., Maddox, S. J. & Sutherland, W. J., 1992, *ApJ Lett.*, **390**, L1.
- Dalton, G. B., Efstathiou, G., Maddox, S. J. & Sutherland, W. J., 1994a, *MNRAS*, **269**, 151.
- Dalton, G. B., Croft, R. A. C., Efstathiou, G., Sutherland, W. J., Maddox, S. J. & Davis, M., 1994b, *MNRAS*, **271**, L47.
- Dalton, G. B., Efstathiou, G., Maddox, S. J. & Sutherland, W. J., 1997, *MNRAS*, **289**, 263.
- Davis, M. & Peebles, P. J. E., 1977, *ApJ*, **34**, 425.
- Diaferio, A. & Geller, M. J., 1997, *ApJ*, **481**, 633.
- Efstathiou, G., Davis, M., Frenk, C. S., & White, S.D. M., 1985, *Ap. J. Suppl.*, **57**, 241.
- Efstathiou, G., 1988, In: *Comets to Cosmology*, 312, ed. Lawrence, A., Springer Verlag.
- Efstathiou, G., 1993, In: *Proceedings of the U.S. National Academy of Sciences*, 4859, ed. 90.
- Eke, V. R., Cole, S. & Frenk, C. S., 1996, *MNRAS*, **282**, 263.
- Gaztañaga, E., 1995, *ApJ*, **454**, 561.
- Gunn, J., Weinberg, D. H., 1995, in *Wide-Field Spectroscopy and the Distant Universe*, eds S.J. Maddox and A. Aragón-Salamanca, World Scientific, Singapore.
- Hatton, S. J. & Cole, S., 1997, *MNRAS*, *submitted*, astro-ph/9707186.
- Hoffman, Y., 1986, *ApJ*, **308**, 493.
- Jenkins, A., Frenk, C. S., Thomas, P. A., Colberg, J. M., White, S.D. M., Couchman, H.M. P., Peacock, J. A., Efstathiou, G., Nelson, A. H., 1998, *ApJ*, *in press*, astro-ph/9709010.
- Kaiser, N., 1987, *Mon. Not. R. astr. Soc.*, **227**, 1.
- Klypin, A., Holtzman, J., Primack, J. & Regös, E., 1993, *ApJ*, **416**, 1.
- Lahav, O., Rees, M. J., Lilje, P. B., Primack, J. R., 1991, *Mon. Not. R. astr. Soc.*, **251**, 128.
- Lilje, P. B. & Efstathiou, G., 1988, *MNRAS*, **231**, 635.
- Lilje, P. B. & Efstathiou, G., 1989, *MNRAS*, **236**, 851.
- Lilje, P. B. & Lahav, O., 1991, *ApJ*, **374**, 29.
- Loveday, J., 1990, *PhD thesis*, Cambridge University.

- Loveday, J., Peterson, B. A., Efstathiou, G., & Maddox, S. J., 1992a, *Ap. J. Lett.*, **390**, 338.
- Loveday, J., Efstathiou, G., Peterson, B. A. & Maddox, S. J., 1992b, *Ap. J. Lett.*, **400**, L43.
- Loveday, J., Efstathiou, G., Maddox, S. J. & Peterson, B. A., 1996, *ApJ*, **468**, 1.
- Maddox, S. J., Efstathiou, G., Sutherland, W. J., & Loveday, J., 1990a, *MNRAS*, **242**, 43.
- Maddox, S. J., Efstathiou, G., Sutherland, W. J., & Loveday, J., 1990b, *MNRAS*, **243**, 692.
- Maddox, S. J., Efstathiou, G., Sutherland, W. J. 1996, *MNRAS*, **283**, 1227.
- Merchán, M. E., Abadi, M. G., Lambas, D. G., & Valotto, C., 1997, *ApJ*, *in press*.
- Mo, H. J., Peacock, J. A., & Xia, X. Y., 1993, *MNRAS*, **260**, 121.
- Moore, B., Frenk, C. S., Efstathiou, G. & Saunders, W., 1994, *MNRAS*, **269**, 742.
- Navarro, J. F., Frenk, C. S., & White, S. D. M., 1997, *ApJ*, *in press*, astro-ph/9611107.
- Peebles, P. J. E., 1980, *The Large-Scale Structure of the Universe*, Princeton University Press.
- Peebles, P. J. E., 1993, *Principles of Physical Cosmology*, Princeton University Press.
- Ratcliffe, A., Shanks, T., Parker, Q. A., Fong, R., 1997, *MNRAS*, *in press*, astro-ph/9702227.
- Reg'os, E. & Geller, M. J., 1989, *Astron. J.*, **98**, 755.
- Saunders, W., Rowan-Robinson, M. & Lawrence, A., 1992, *MNRAS*, **258**, 134.
- Schectman, S. A., Landy, S. D., Oemler, A., Tucker, D. L., Lin, H., Kirshner, R. P., Schecter, P. L., 1996, **470**, 172.
- Seldner, M. & Peebles, P. J. E., 1977, *ApJ*, **215**, 703.
- Summer, F. J., Davis, M. & Evrard, A., 1995, *ApJ*, **454**, 1.
- van Haarlem, M., 1992, *PhD thesis*, Leiden University.
- van Haarlem, M., Frenk, C. S. & White, S. D. M., 1997, *MNRAS*, **287**, 817.
- van Haarlem, M., Cayon, L., de la Cruz, C. G., Martinez-Gonzalez, E. & Rebolo, R., 1993, *MNRAS*, **264**, 71.
- van Haarlem, M. & van de Weygaert, R., 1993, *ApJ*, **308**, 499.
- Villumsen, J. V. & Davis, M., 1986, *ApJ*, **308**, 499.
- Weinberg, D. H., in *Wide-Field Spectroscopy and the Distant Universe*, eds. S. J. Maddox and A. Aragón-Salamanca, (Singapore: World Scientific), 129, astro-ph/9409094.
- White, S. D. M., Efstathiou, G. & Frenk, C. S., 1993, *MNRAS*, **262**, 1023.
- Wright, E. L., Smoot, G. F., Kogut, A., Hinshaw, G., Teno-río, L., Lineweaver, C., Bennett, C. L. & Lubin, P. M., 1994, *ApJ*, **420**, 1.
- Yahil, A., Sandage, A. & Tammann, G. A., 1980, *ApJ*, **242**, 448.
- Yahil, A., 1985. In: *The Virgo Cluster of Galaxies*, 359, eds Richter, O. G. & Binggeli, B., ESO : Garching.